

Dinâmica Neuro-Olfativa: perspectivas eletroencefalográficas de disfunção

Artigo Original

Autores

Francisco Alves de Sousa

Centro Hospitalar Universitário do Porto

Sara Costa

Centro Hospitalar Universitário do Porto

João Tavares Correia

Centro Hospitalar Universitário do Porto

Sara Azevedo

Centro Hospitalar Universitário do Porto

Ana Nóbrega Pinto

Centro Hospitalar Universitário do Porto

Marilene Santos

Centro Hospitalar Universitário do Porto

Luís Meireles

Centro Hospitalar Universitário do Porto

Resumo

Introdução: O olfato relaciona-se com cognição. Compreender a atividade cerebral em contexto de disfunção olfativa (DO) poderia alargar o conhecimento sobre o olfato e os tratamentos associados, como o treino olfativo (TO).

Objetivos: Investigar como a DO afeta a atividade cerebral em resposta a aromas.

Métodos: Para realizar este estudo prospetivo, foi recrutado um grupo de indivíduos com DO e controlos saudáveis. Os participantes inalaram aromas usados no TO e a atividade cerebral foi medida através de Eletroencefalografia (EEG).

Resultados: A ativação cerebral diferiu entre os grupos para 3 dos 4 aromas utilizados. Rosa: menor ativação Alfa 1 no grupo DO ($p=0,021$); Eucalipto: menor ativação Beta 1 na DO ($p = 0,037$); Cravinho: menor ativação Gama 1, Beta 2 e Alfa 2 no grupo DO ($p < 0,05$); atividade Delta mais alta no grupo DO ($p = 0,019$). Não foram encontradas diferenças entre grupos para a inalação de limão.

Conclusões: Este estudo explorou a atividade cerebral durante a perceção olfativa na DO. A personalização do TO com base no aroma e na resposta cerebral individual apresenta-se como uma abordagem promissora. É necessária mais investigação.

Palavras-chave: Olfato; Disfunção Olfativa; Treino Olfativo; Eletroencefalografia; Atividade Cerebral; Aroma/Odor; Efeitos Neurocognitivos; Reabilitação

Introduction

Olfactory pathways extend far beyond the simple detection of odors, acting as a vital link between the sense of smell and various brain functions, including cognition, behavior and immune system activity^{1,2}. This unique relationship between smell and cognition is further underscored by the evolutionary development of the olfactory system. Unlike other sensory modalities, olfaction bypasses the thalamus, a relay center for most senses, and projects directly to the limbic system and other higher brain regions^{2,3}. This intimate connection suggests that odors can trigger

Correspondência:

Francisco Alves de Sousa
franciscoalvesousa@gmail.com

Artigo recebido a 8 de Abril de 2024.
Aceite para publicação a 7 de Junho de 2024.

immediate emotional responses and influence behavior⁴. The interplay between olfaction and the limbic system, integrating emotions and memory, is likely to be a key factor in the success of OT. In fact, studies suggest a strong connection between olfactory training (OT) and cognition⁵. Positron emission tomography (PET) studies further support the role of the orbitofrontal cortex in processing emotionally charged odors and decision-making^{2,3,6}. Ultimately, Electroencephalographic (EEG) readings can be performed at this region to detect activity in response to odors⁷. With this in mind, the primary objective of this work was to investigate the neurocognitive effects of OT in individuals with OD compared to healthy controls, by employing EEG to assess brain activity during the inhalation of common OT scents. Secondary objectives included examining potential relationships between specific odorants and unique brain activation patterns measured by EEG, as well as investigating whether individual variations in baseline smell function influenced the response to olfactory training as measured by brain activity.

Material and Methods

Participants

This case-control prospective study recruited participants from two groups: individuals diagnosed with olfactory dysfunction (OD) and healthy controls with normal olfactory function. To confirm or rule out OD, all participants underwent a standardized olfactory perception test (OPT) using the Sniffin' Sticks threshold test⁸. A cut-off OPT of 7 was chosen to define hyposmia accordingly to relevant Literature and manufacturers' information⁹⁻¹². Exclusion criteria for controls were: age < 18 years, OPT ≤ 7, chronic rhinosinusitis, history of head trauma with loss of consciousness, documented pre-existing olfactory dysfunction, pregnancy, previous neurosurgery or endoscopic nasal surgery, known olfactory bulb lesions on imaging, known neurologic disease (Parkinson's, Dementia, Epilepsy). Inclusion criteria for cases: age > 18 years, reporting OD with objective OPT ≤ 7.

Experimental Environment

The experiment was conducted in a controlled environment designed to minimize external influences on brain activity measured by EEG. The room was maintained at a standard temperature and ensured to be quiet and well-ventilated to optimize participant comfort and focus during the testing session.

Experimental Protocol

The experiment consisted of a series of odor exposures while EEG data was collected. Each participant underwent the following procedures:

1. Baseline Measurement:

The session began with a one-minute baseline recording where participants were exposed only to clean, odorless distilled water. This established EEG resting brain activity before scent introduction. To ensure a neutral starting point for each odor presentation, this baseline measurement was repeated between each scent inhalation. Participants then proceeded with the one-minute exposure to the designated odor.

2. Odor Presentation:

Following the baseline, participants were exposed to four different odorants: rose, lemon, eucalyptus, and clove. Each odor was presented for a duration of one minute. During odor presentation, participants were instructed to stay quiet and inhale deeply while focusing on the specific scent, minimizing any mental distractions.

3. Inter-Stimulus Interval:

A brief inter-stimulus interval was included between each baseline-odor presentation sequence (approximately 30 seconds). This interval allowed participants' brains to return to a relatively neutral state.

EEG Recording and Analysis

This study employed a single-channel electroencephalography (EEG) sensor to record brain activity during odor exposure.

The sensor, a NeuroSky Mindwave 2, was positioned on the participant's scalp according to the international 10-20 system at the AF3 channel, a location associated with olfactory processing. This non-invasive sensor offers a sampling rate of 512 Hz, capturing EEG data at a rate of approximately once per second. The EEG data was analyzed to assess the power spectrum frequencies of various brain waves, including Theta (θ), Delta (δ), Alpha 1 (α_1), Alpha 2 (α_2), Beta 1 (β_1), Beta 2 (β_2), and Gamma (γ) waves. Each of these wave types is associated with distinct brain states. For example, Alpha waves are typically associated with relaxed wakefulness, while Beta waves are linked to focused attention (see Table 1). The performance-to-baseline ratios for each brain-wave/odor sequence were recorded in a database. Performance-to-baseline ratios: refers to the ratio of the EEG activity during odor exposure (performance) to the EEG activity during the baseline recording (baseline). This ratio provides a measure of how odor exposure affects brain activity

compared to a resting state. To provide a more comprehensive understanding of brain activity during odor exposure, an "Activation Index" (AI) was calculated. This index aimed to capture the balance between mental activation and resting/sleep states. The AI was derived by dividing the sum of the power spectrum values for activating waves (Gamma 1, Gamma 2, Beta 1, and Beta 2) by the sum of the power spectrum values for resting/sleep waves (Theta, Delta, Alpha 1, and Alpha 2). A higher AI score indicates a greater level of mental activation during odor exposure.

Ethics

Informed consent was obtained for all patients and was submitted to local Ethics Committee approval. Design complies with the Declaration of Helsinki's ethical standards.

Data Analysis

The collected EEG data, including individual wave power spectrums and the calculated AI scores, were statistically analyzed.

Table 1
Different types of brain waves and respective mental state associations

Brainwave Type	Frequency (Hz)	Function	Association	Image
Delta (δ)	0.5 - 4 Hz	Deep sleep, unconscious processing	Deep relaxation, unconscious processing	
Theta (θ)	4 - 8 Hz Theta 1 (4-5 Hz) Theta 2 (5-7 Hz) Theta 3 (7-8 Hz)	Daydreaming, meditation, drowsiness	Daydreaming, meditation, drowsiness	
Alpha (α)	8 - 12 Hz Alpha 1 (8-10 Hz) Alpha 2 (10-12 Hz)	Relaxed wakefulness, calmness	Relaxed wakefulness, calmness	
Beta (β)	13 - 30 Hz Beta 1 (13-16 Hz) Beta 2 (16-20 Hz) Beta 3 (20-30 Hz)	Focused attention, alertness	Focused attention, alertness	
Gamma (γ)	30 - 80 Hz Gamma 1 (30-40 Hz) Gamma 2 (40-50 Hz) Gamma 3 (50-80 Hz)	Higher-order processing, information binding	Higher-order processing, information binding	

This analysis compared brain activity patterns between the OD and control groups for each odorant presented. The goal was to identify any significant differences in neural responses to the various scents between individuals with and without OD. Statistical analyses were conducted using IBM SPSS Statistics 29 software. Descriptive statistics were employed, presenting categorical variables as percentages and continuous variables as means and standard deviations. The Shapiro-Wilk test confirmed normal distribution for continuous variables. Bivariate analysis explored asymmetries in measurements between groups by means of independent t-test. Pearson Chi-square/ Fisher's tests (95% confidence intervals) were used for categories and Spearman's test for continuous variables. Mixed ANOVAs were performed to compare brain wave responses to each odor (test for between-subjects effects being group). An additional mixed ANOVA with within-subjects factor being type of odor and between-subjects factor being group (OD or control) checked asymmetries in activation index. This allowed the production of graphical representations on differences between groups for each scent. P-values derived from ANOVA models were

obtained after Mauchly's test of sphericity. When sphericity assumption was violated, Greenhouse-Geisser test was preferred. All reported p-values are two-tailed, with a significance level set at $p \leq 0.05$.

Results

Population

A general description of the main variables is displayed on Table 1. A total of 34 patients were enrolled, 15 cases and 19 controls. 21% of cases and 46.7% of controls were male. No differences were noted in age and sex distribution between groups ($p > 0.05$). Within the 15 participants diagnosed with OD, nine (60%) had post-viral etiology, three (20%) had chronic rhinosinusitis, two (13.3%) were classified as idiopathic, and one (6.7%) post-traumatic. A within-group analysis revealed no significant effect of sex on the activation index for any of the tested odors (rose, lemon, eucalyptus, clove) in either the olfactory dysfunction (OD) group or the control group (all $p > 0.05$). This suggests that sex does not play a significant role in modulating the neural responses to these specific odors within the context of this study. Figure 1 displays mean activation index for each OD etiology.

Figure 1

Average activation index for each odorant, stratified by the underlying cause of olfactory dysfunction




Table 1
Average activation index for each odorant, stratified by the underlying cause of olfactory dysfunction

Odor	Feature	Cases	Controls	p-value	
	Sample Size (n)	15	19	-	
	Sex Distribution	7 M / 8 F	4 M / 15 F	0.113	
	Age (years)	35 ± 11	37 ± 10	0.588	
	Olfactory perception threshold	9.94 ± 2.23	1.33 ± 1.50	< 0.001	
Rose	(Performance/ baseline) ratio	Delta (δ)	1.10 ± 0.23	1.03 ± 0.20	0.346
		Theta (θ)	0.97 ± 0.07	1 ± 0.08	0.272
		Alpha 1 (α_1)	0.90 ± 0.13	1.05 ± 0.19	0.021
		Alpha 2 (α_2)	0.91 ± 0.19	0.98 ± 0.19	0.306
		Beta 1 (β_1)	0.93 ± 0.16	1.04 ± 0.18	0.073
		Beta 2 (β_2)	0.97 ± 0.21	1.04 ± 0.17	0.336
		Gamma 1 (γ_1)	1.01 ± 0.25	1.10 ± 0.21	0.302
	Gamma 2 (γ_2)	0.93 ± 0.27	1.07 ± 0.27	0.149	
	Activation index	0.98 ± 0.19	1.05 ± 0.16	0.340	
Lemon	(Performance/ baseline) ratio	Delta (δ)	0.98 ± 0.17	1.02 ± 0.20	0.482
		Theta (θ)	1.05 ± 0.12	0.99 ± 0.13	0.149
		Alpha 1 (α_1)	1.03 ± 0.15	1.04 ± 0.19	0.887
		Alpha 2 (α_2)	1.02 ± 0.15	1.03 ± 0.31	0.847
		Beta 1 (β_1)	1.05 ± 0.16	1.03 ± 0.20	0.723
		Beta 2 (β_2)	1.05 ± 0.19	1.04 ± 0.23	0.863
		Gamma 1 (γ_1)	1.06 ± 0.22	1.08 ± 0.26	0.872
	Gamma 2 (γ_2)	1.05 ± 0.21	1.03 ± 0.21	0.796	
	Activation index	1.04 ± 0.16	1.02 ± 0.14	0.767	
Eucalyptus	(Performance/ baseline) ratio	Delta (δ)	1.14 ± 0.21	1.08 ± 0.22	0.421
		Theta (θ)	0.99 ± 0.09	1.03 ± 0.10	0.290
		Alpha 1 (α_1)	0.91 ± 0.14	0.97 ± 0.20	0.309
		Alpha 2 (α_2)	0.88 ± 0.16	0.98 ± 0.17	0.108
		Beta 1 (β_1)	0.88 ± 0.14	0.99 ± 0.15	0.037
		Beta 2 (β_2)	0.92 ± 0.18	0.98 ± 0.21	0.379
		Gamma 1 (γ_1)	1 ± 0.18	1.03 ± 0.20	0.654
	Gamma 2 (γ_2)	0.93 ± 0.21	0.90 ± 0.23	0.687	
	Activation index	0.95 ± 0.16	0.96 ± 0.16	0.872	
Clove	(Performance/ baseline) ratio	Delta (δ)	1.12 ± 0.21	0.97 ± 0.10	0.019
		Theta (θ)	1 ± 0.06	1.02 ± 0.09	0.472
		Alpha 1 (α_1)	0.95 ± 0.16	0.96 ± 0.10	0.656
		Alpha 2 (α_2)	0.92 ± 0.18	1.05 ± 0.15	0.039
		Beta 1 (β_1)	0.95 ± 0.17	1.01 ± 0.14	0.261
		Beta 2 (β_2)	0.96 ± 0.16	1.09 ± 0.13	0.019
		Gamma 1 (γ_1)	1.02 ± 0.12	1.13 ± 0.18	0.046
	Gamma 2 (γ_2)	0.95 ± 0.15	1.03 ± 0.16	0.100	
	Activation index	0.97 ± 0.12	1.07 ± 0.12	0.044	

M = Male, F = Female; Values are presented as Mean ± Standard Deviation; p-value indicates statistical significance between cases and controls. A value of less than 0.05 (in bold) is considered statistically significant.

Bivariate analysis

Rose inhalation showed a significant difference between OD and control groups, with lower Alpha 1 activation (OD: 0.90 ± 0.13 vs controls: 1.05 ± 0.19 , $p=0.021$). Eucalyptus inhalation revealed lower Beta 1 activation in OD patients (OD: 0.88 ± 0.14 vs controls: 0.99 ± 0.15 , $p = 0.037$). Clove inhalation presented significant differences in various frequency bands: Gamma 1, Beta 2 and Alpha 2 were lower in OD (OD $\gamma 1$: 1.01 ± 0.12 vs controls $\gamma 1$: 1.13 ± 0.18 , $p = 0.046$; OD $\beta 2$: 0.96 ± 0.16 vs controls $\beta 2$: 1.10 ± 0.13 , $p = 0.019$; OD $\alpha 2$: 0.92 ± 0.17 vs controls $\alpha 2$: 1.05 ± 0.15 , $p = 0.039$); and Delta activity was higher in OD for clove inhalation (OD δ : 1.12 ± 0.21 vs controls δ : 0.97 ± 0.1 , $p = 0.019$). The Activation Index significantly differed during clove inhalation, being lower in OD (OD AI: 0.97 ± 0.11 vs controls: 1.1 ± 0.12 , $p = 0.044$). Remarkably, no differences were found in any parameter related to lemon inhalation ($p > 0.05$). There was a positive correlation between baseline OPT and activation index for clove ($p = 0.038$), but not for any other scent ($p = 0.187$ for rose; $p = 0.616$ for lemon; $p = 0.320$ for eucalyptus).

Multivariate analysis

Additional ANOVA models were created in order to further validate the findings from the former analysis (Table 2).

Brain wave responses to Rose: OD vs controls

The main effect of olfactory dysfunction on brain activation to rose was not significant in this model ($F(1,59) = 3.076$, $p = 0.090$). Nevertheless, there was a tendency for significance (see Figure 2). Analysis of performance-to-baseline ratios across frequency bands revealed no statistically significant differences ($F(7, 203) = 1.748$, $p = 0.182$).

Brain wave responses to Lemon: OD vs controls

The main effect of olfactory dysfunction on brain activation to lemon was not significant in this model ($F(1,29) = 0.010$, $p = 0.922$; see Figure 3). Analysis of performance-to-baseline ratios across frequency bands revealed no statistically significant differences ($F(7, 203) = 0.523$, $p = 0.620$).

Figure 2
Mean brain responses (performance/baseline ratio) to rose: OD vs controls




Figure 3
Mean brain responses (performance/baseline ratio) to lemon: OD vs controls




Figure 4
Mean brain responses (performance/baseline ratio) to eucalyptus: OD vs controls




Figure 5
Mean brain responses (performance/baseline ratio) to clove: OD vs controls





Figure 6
Mean activation index for each odor inhalation: OD vs controls



Brain wave responses to Eucalyptus: OD vs controls

The main effect of olfactory dysfunction on brain activation to eucalyptus was not

significant in this model ($F(1,28) = 1.346, p = 0.256$; see Figure 4). Analysis of performance-to-baseline ratios across frequency bands revealed statistically significant differences

($F(7, 196) = 4.727, p = 0.007$), since delta wave signal clearly predominated in both OD patients and controls (see Figure 4).

Brain wave responses to Clove: OD vs controls

The main effect of olfactory dysfunction on brain activation to eucalyptus was not significant in this model ($F(1,28) = 2.905, p = 0.099$). Nevertheless, there was a tendency for significance (see Figure 5). Analysis of performance-to-baseline ratios across frequency bands revealed a statistical tendency towards significance ($F(7, 196) = 2.777, p = 0.058$) (see Figure 5).

Activation index: OD vs controls

The overall main effect of olfactory dysfunction on activation index to eucalyptus was not significant in this model ($F(1,27) = 2.465, p = 0.128$). Nevertheless, there was a marked difference concerning clove as showed in the bivariate analysis (see Table 2 and Figure 6).

Discussion

This study successfully achieved its central aim: investigating the neurocognitive effects of OD on brain activity during odor exposure using EEG. While the primary focus compared responses between individuals with OD and healthy controls, the broader implications of these findings extend to the potential for personalized olfactory training (OT) regimens in the future. The one-minute odor inhalation period was deliberately chosen to mirror a typical olfactory training session. This approach aimed to simulate the repeated exposure to specific scents employed in established OT protocols, enhancing the generalizability of our findings to real-world OT applications.

The analysis of EEG data revealed intriguing insights into how participants processed odors. Notably, the rose inhalation task yielded a significant difference in Alpha 1 wave activation between the groups, with the control group exhibiting a stronger response (Table 2). This finding, potentially linked to a state of relaxed wakefulness during rose odor processing, suggests that OD may alter brain

activity in this setting. Similarly, patients with OD exhibited a lower Beta 1 signal during eucalyptus inhalation compared to the control group. Beta waves are associated with focused attention (7) (see Table 1), suggesting that individuals with OD may have reduced frontal cortex response when encountering the eucalyptus odor. Clove exposure yielded the most striking results. OD patients showed lower activation in specific bands (Gamma 1, Beta 2, Alpha 2) compared to controls. This suggests potentially poorer neural processing for higher cognitive functions (attention, memory – related to Gamma and Beta activity) and possibly reduced relaxation (Alpha 2) in the OD group during clove inhalation. Additionally, higher Delta activity (deep sleep/inactive state) and a lower Activation Index (overall engagement) in the OD group further support this notion of diminished neural response to clove. Analysis using ANOVA revealed high delta wave activity in both OD and control groups during eucalyptus inhalation (Figure 4). Delta waves are typically associated with states of deep relaxation or sleep⁷. This finding aligns with research demonstrating the anxiolytic (anxiety-reducing) effects of 1,8-cineole, a key component of eucalyptus oil¹³. ANOVA analyses of other odors (excluding lemon) revealed trends towards significance, suggesting potential odor-specific effects. However, statistical power may have been limited by sample size, so a larger population is warranted to confirm these trends and definitively identify meaningful changes in neural responses. Interestingly, and unlike other odors in this study, lemon inhalation did not yield significant differences in brain activity patterns between OD and control groups. While initially surprising, this finding could be explained by the unique properties of lemon, as it possesses a powerful odor profile, with high concentration of volatile compounds¹⁴. These compounds are well-established olfactory stimulants, and research suggests they can even trigger the release of the stress hormone epinephrine¹⁵. This potent activation, bypassing potential olfactory deficits in

the OD group, could explain the observed similarity in brain activity patterns between both groups. The evocative power of scent, captured so vividly by Marcel Proust in his novel "À la recherche du temps perdu", has long fascinated researchers^{12,13}. This phenomenon, now known as the Proust Phenomenon, reflects the unique ability of odors to trigger vivid and often emotional autobiographical memories¹⁴. The interconnexion between olfaction and the limbic system (orbitofrontal cortex, insula, anterior cingulate cortex, amygdala and hippocampus) explains the inseparable relation between odors and memories/emotions¹⁵⁻¹⁷. This "higher olfactory functions" play a broader role in shaping human behavior beyond just basic odor detection^{2,3,18-20}. Considering OT, individual differences in both olfactory function and the emotional connections to specific scents could significantly impact how individuals respond to the training odors¹⁵. These variations can ultimately influence the effectiveness of the training program²⁵. Therefore, personalizing OT by identifying the most effective scents for each individual holds immense promise for optimizing treatment outcomes. This study lays the groundwork for this crucial tailoring approach by demonstrating the potential for specific odor stimuli to elicit distinct neural responses. The process of chemosensory perception involves distinct neural networks: the olfactory network, responsible for odor detection and identification; the somatosensory network, mediating the perception of trigeminal sensations such as pungency or coolness; and the integrative network, responsible for combining and interpreting olfactory and trigeminal information to create a unified percept²⁶. Understanding the functional connectivity within and between these networks is crucial for comprehending how the brain processes and interprets chemosensory stimuli. Our study, while primarily focused on EEG correlates of odor perception, highlights the need for future research into the functional connectivity of these networks. Advanced neuroimaging techniques like functional magnetic resonance imaging (fMRI), combined

with EEG, could elucidate how the olfactory, somatosensory, and integrative networks interact during odor exposure, potentially unveiling individual differences in odor perception and the potential for personalized OT interventions²⁷. There are limitations to acknowledge. Primarily, the sample size was relatively small, potentially impacting the generalizability of the findings. Only OPT were measured in the psychophysical domain, so that identification and discrimination were not evaluated in this sample. This choice was made as OPT are considered a fundamental indicator of peripheral olfactory function, whereas identification and discrimination are believed to be influenced by more complex olfactory and cognitive processes^{11,28,29}. Additionally, employing multi-channel EEG systems would offer a more comprehensive picture of brain activity during odor exposure. Future studies incorporating multi-channel EEG could provide a more comprehensive understanding of the specific brain regions and networks involved in odor perception and potentially in response to olfactory training. The inclusion of diverse etiologies of olfactory dysfunction in this study may also have introduced variability in the results, potentially masking etiology-specific patterns of brain activation. We recognize that the design of this study limits our ability to draw definitive conclusions about the long-term effects of OT. Future studies with extended follow-up periods are therefore necessary. We also acknowledge the importance of considering the interplay between the olfactory, somatosensory, and integrative networks in olfactory processing. Incorporating functional connectivity analyses could shed light on how these networks interact to shape odor perception and how OT may modulate their connectivity.

Conclusion

This study investigated the impact of OD on brain activity during odor exposure using EEG, highlighting the multifaceted nature of olfaction. Findings revealed differences in central brain integration between individuals

with OD and healthy controls. These observations, coupled with the known link between olfaction, memory, and emotion, suggest promising avenues for OT. Future research could explore individual variations in smell perception and corresponding brain responses, in view of personalizing OT protocols. Additionally, exploring the effects of different odor profiles and incorporating advanced neuroimaging could further optimize OT regimens and unravel OD neural mechanisms. Ultimately, this deeper understanding can lead to more effective interventions for OD patients.

Acknowledgments

We extend our sincere gratitude to all the clinical and non-clinical staff of the Otorhinolaryngology Department for endorsing this work.

Conflict of Interests

The authors declare that they have no conflict of interest regarding this article.

Data Confidentiality

The authors declare that they followed the protocols of their work in publishing patient data.

Human and animal protection

The authors declare that the procedures followed are in accordance with the regulations established by the directors of the Commission for Clinical Research and Ethics and in accordance with the Declaration of Helsinki of the World Medical Association.

Privacy policy, informed consent and Ethics committee authorization

All the processed data were based in published reports that fulfilled privacy policy and ethical considerations.

Financial support

This work did not receive any grant contribution, funding or scholarship.

Scientific data availability

There are no publicly available datasets related to this work.

References

1. Strous RD, Shoenfeld Y. To smell the immune system: olfaction, autoimmunity and brain involvement. *Autoimmun Rev.* 2006 Nov;6(1):54-60. doi: 10.1016/j.autrev.2006.07.002.
2. Savic I. Imaging of brain activation by odorants in humans. *Curr Opin Neurobiol.* 2002 Aug;12(4):455-61. doi: 10.1016/s0959-4388(02)00346-x.
3. Bastir M, Rosas A, Gunz P, Peña-Melian A, Manzi G, Harvati K. et al. Evolution of the base of the brain in highly encephalized human species. *Nat Commun.* 2011 Dec 13;2:588. doi: 10.1038/ncomms1593.
4. Kivity S, Ortega-Hernandez OD, Shoenfeld Y. Olfaction--a window to the mind. *Isr Med Assoc J.* 2009 Apr;11(4):238-43.
5. Vance DE, Del Bene VA, Kamath V, Frank JS, Billings R, Cho DY. et al. Does olfactory training improve brain function and cognition? A systematic review. *Neuropsychol Rev.* 2024 Mar;34(1):155-191. doi: 10.1007/s11065-022-09573-0.
6. Bechara A, Damasio H, Damasio AR. Emotion, decision Making and the orbitofrontal cortex. *Cereb Cortex.* 2000 Mar;10(3):295-307. doi: 10.1093/cercor/10.3.295.
7. Laohakangvalvit T, Sripian P, Nakagawa Y, Feng C, Tazawa T, Sakai S. et al. Study on the psychological states of olfactory stimuli using electroencephalography and heart rate variability. *Sensors (Basel).* 2023 Apr 16;23(8):4026. doi: 10.3390/s23084026.
8. Denzer MY, Gailer S, Kern DW, Schumm LP, Thuerauf N, Kornhuber J. et al. Quantitative Validation of the n-Butanol Sniffin' Sticks Threshold Pens. *Chemosens Percept.* 2014;7(2):91-101. doi: 10.1007/s12078-014-9168-1
9. Ribeiro JC, Simões J, Silva F, Silva ED, Hummel C, Hummel T. et al. Cultural adaptation of the portuguese version of the "Sniffin' Sticks" Smell Test: reliability, validity and normative data. *PLoS One.* 2016 Feb 10;11(2):e0148937. doi: 10.1371/journal.pone.0148937.
10. Hsieh JW, Keller A, Wong M, Jiang R-S, Vosshall LB. SMELL-S and SMELL-R: olfactory tests not influenced by odor-specific insensitivity or prior olfactory experience. *Proc Natl Acad Sci U S A.* 2017 Oct 24;114(43):11275-11284. doi: 10.1073/pnas.1711415114
11. Hummel T, Kobal G, Gudziol H, Mackay-Sim A. Normative data for the "Sniffin' Sticks" including tests of odor identification, odor discrimination, and olfactory thresholds: an upgrade based on a group of more than 3,000 subjects. *Eur Arch Otorhinolaryngol.* 2007 Mar;264(3):237-43. doi: 10.1007/s00405-006-0173-0.
12. Oleszkiewicz A, Schriever VA, Croy I, Hähner A, Hummel T. Updated Sniffin' Sticks normative data based on an extended sample of 9139 subjects. *Eur Arch Otorhinolaryngol.* 2019 Mar;276(3):719-728. doi: 10.1007/s00405-018-5248-1.
13. Kim KY, Seo HJ, Min SS, Park M, Seol GH. The effect of 1,8-cineole inhalation on preoperative anxiety: a randomized clinical trial. *Evid Based Complement Alternat Med.* 2014;2014:820126. doi: 10.1155/2014/820126.
14. Kiecolt-Glaser JK, Graham JE, Malarkey WB, Porter K, Lemeshow S, Glaser R. Olfactory influences on mood

- and autonomic, endocrine, and immune function. *Psychoneuroendocrinology*. 2008 Apr;33(3):328-39. doi: 10.1016/j.psyneuen.2007.11.015.
15. Martial C, Poirrier AL, Pottier L, Cassol H, Mortaheb S, Panda R. et al. From nose to brain: the effect of lemon inhalation observed by whole brain voxel to voxel functional connectivity. *Cortex*. 2023 Aug;165:119-128. doi: 10.1016/j.cortex.2023.04.012.
16. Chu S, Downes JJ. Proust nose best: odors are better cues of autobiographical memory. *Mem Cognit*. 2002 Jun;30(4):511-8. doi: 10.3758/bf03194952.
17. Jellinek JS. Proust remembered: has Proust's account of odor-cued autobiographical memory recall really been investigated? *Chem Senses*. 2004 Jun;29(5):455-8; author reply 459-61. doi: 10.1093/chemse/bjh043.
18. Herz RS. The role of odor-evoked memory in psychological and physiological health. *Brain Sci*. 2016 Jul 19;6(3):22. doi: 10.3390/brainsci6030022.
19. Kadohisa M. Effects of odor on emotion, with implications. *Front Syst Neurosci*. 2013 Oct 10;7:66. doi: 10.3389/fnsys.2013.00066.
20. Soudry Y, Lemogne C, Malinvaud D, Consoli S-M, Bonfils P. Olfactory system and emotion: common substrates. *Eur Ann Otorhinolaryngol Head Neck Dis*. 2011 Jan;128(1):18-23. doi: 10.1016/j.anorl.2010.09.007.
21. Gottfried JA, Smith APR, Rugg MD, Dolan RJ. Remembrance of odors past: human olfactory cortex in cross-modal recognition memory. *Neuron*. 2004 May 27;42(4):687-95. doi: 10.1016/s0896-6273(04)00270-3
22. Fagundo AB, Jiménez-Murcia S, Giner-Bartolomé C, Islam MA, de la Torre R, Pastor A. et al. Modulation of higher-order olfaction components on executive functions in humans. *PLoS One*. 2015 Jun 17;10(6):e0130319. doi: 10.1371/journal.pone.0130319
23. Pieniak M, Rokosz M, Ivcevic Z, Reichert A, Żyżelewicz B, Nawrocka P. et al. Olfactory training improves emotion matching ability in 6–9 years old children — preliminary evidence. *J Sens Stud* 2024 Apr 1;39(2):e12912. doi.org/10.1111/joss.12912.
24. Okuni I, Ebihara S. Intensive olfactory training and emotional memory in patients with dementia. *Geriatr Gerontol Int*. 2022 Feb;22(2):185-186. doi: 10.1111/ggi.14344.
25. Pieniak M, Oleszkiewicz A, Avaro V, Calegari F, Hummel T. Olfactory training – thirteen years of research reviewed. *Neurosci Biobehav Rev*. 2022 Oct;141:104853. doi: 10.1016/j.neubiorev.2022.104853.
26. Small DM. Flavor is in the brain. *Physiol Behav*. 2012 Nov 5;107(4):540-52. doi: 10.1016/j.physbeh.2012.04.011.
27. Seubert J, Ohla K, Yokomukai Y, Kellermann T, Lundström JN. Superadditive opercular activation to food flavor is mediated by enhanced temporal and limbic coupling. *Hum Brain Mapp*. 2015 May;36(5):1662-76. doi: 10.1002/hbm.22728.
28. Kollndorfer K, Kowalczyk K, Hoche E, Mueller CA, Pollak M, Trattng S. et al. Recovery of olfactory function induces neuroplasticity effects in patients with smell loss. *Neural Plast*. 2014;2014:140419. doi: 10.1155/2014/140419.
29. Hedner M, Larsson M, Arnold N, Zucco GM, Hummel T. Cognitive factors in odor detection, odor discrimination, and odor identification tasks. *J Clin Exp Neuropsychol*. 2010 Dec;32(10):1062-7. doi: 10.1080/13803391003683070.